



Bayesian Analysis of Average Age Prevalence of Hypertensive Patients in South Western Part of Nigeria

Olawale Basheer Akanbi

Department of Statistics, University of Ibadan, Ibadan

Email: muhdbashola@yahoo.com

Article history: Received 13 May 2018, Revised 20 June 2018, Accepted 2 July 2018, Published 9 July 2018.

Abstract: Hypertension is a non-communicable disease which is mostly related to age distribution and is rampant in Nigeria. One is likely to develop high blood pressure as long he is getting older. This paper was aimed to determine the average age that one is likely to develop hypertension in Ibadan North Local Government (IBLG). Bayesian Inference was used to analysis this problem. Both uniform and Normal priors were used for the probability update. The month that had the highest number of cases was January and the minimum was recorded in November. It was also observed that 862 (73.4%) of hypertensive patients in the IBLG were female while 312 (26.6%) of them were male. Age group with the highest case was between (52 and 61) years (29.1%) while only one person (0.1%) fell in the age group of 92years and above. The classical estimate, the posterior means for uniform and conjugate priors were (53.5, 53.4 & 53.7) years respectively. The 95% credible interval for the age using conjugate prior was [47.23, 60.17]. It was recommended that an informative normal prior and normal likelihood should be considered for analysing average age prevalence of hypertensive patients in IBLG.

Keywords: Conjugate prior, posterior distribution, Normal distribution, Uniform Distribution, Blood pressure.

AMS Subject Classification: 68M15, 62F15

1. Introduction

Infectious and Non Communicable Diseases (NCDs) are serious issues affecting the health sector. The NCDs Cardiovascular, diabetes, cancer, and chronic respiratory diseases are common challenges to human health and development. These four diseases are the world's biggest killers, causing in excess of 36 million deaths each year of which 80% are in low- and middle- income countries (World Health Organization (WHO) 2011). Unhealthy diet, physical inactivity, tobacco use and the harmful use of alcohol are the behavioral risk factors responsible for about 80% of coronary heart disease and cerebrovascular disease deaths (WHO 2013). The WHO also predicted globally, between 2006 and 2015, NCDs deaths to increase by 17% with highest in the African region by 27%. The most rampant NCDs is hypertension. Hypertension is a common and a major public health problem (Murray et al 1997). It is a serious threat to the health of adults usually in Africa. Hypertension was defined as systolic Blood Pressure (BP) ≥ 140 mmHg or as diastolic BP ≥ 90 mmHg or being on drug therapy. Recently, the prevalence of hypertension was observed to be 28% in North America, 44% in Western Europe (Wolf-Maier et al 2003), 28.3% for crude and 27.3% for age-standardized in urban Accra of Ghana West Africa (Amoah AGB 2003).

Blood pressure is therefore the force of blood against the artery walls as it circulates through the body. Blood pressure is written as two numbers. The first (systolic) number represents the pressure in blood vessels when the heart beats. The second (diastolic) number represents the pressure in the vessels when the heart rests between beats. High blood pressure can cause serious damage to health. It can harden the arteries, decreasing the flow of blood and oxygen to the heart. Perhaps, this reduction of the blood flow can cause chest pain, angina or heart failure which occurs when the heart cannot pump enough blood and oxygen to other organs, heart attack which occurs when the blood supply to the heart is blocked and heart muscle cells die from lack of oxygen. The longer the blood flow is blocked, the greater the damage to the heart. High blood pressure can burst or block arteries that supply blood and oxygen to the brain causing stroke. High blood pressure cannot be cured. It can however, be managed very effectively through lifestyle changes and, when needed medication. In most cases, the cause of high blood pressure is not known. In fact, high blood pressure usually doesn't have symptoms. This is why it is sometimes called "the silent killer." However, there are known risk factors for high blood pressure in which age is included. This deadly disease is not only common among the non-medical staff but also the Healthcare Workers (HCW) especially the nurses. For instance, the report of the Medical Research Council (MRC) of the South Africa showed that NCDs (hypertension) accounted for 37% of deaths (Steyn et al 2006) and recently 20% of deaths were attributed to hypertension among the HCW in South Africa (Skaal L and Pengpid S 2011). This compared with that of the Mexico HCW of 22% of deaths caused by the NCDs (Gonzalez-Valazquez et al 2006).

2. Methodology

Considering the information in the past and also in the literature, hypertension is always traced back to age distributions especially among the old persons. Therefore, suggest the Bayesian procedure in order to improve on the information. The use of Bayesian reasoning in statistics and economics has roughly tripled since 1970 (Poirer, 2006).

Constructing the Likelihood Function

The observation y is a random variable taken from a normal distribution with mean μ and variance σ^2 which is assumed known.

The likelihood function is given by:

$$f(y_i/\theta) = L(\theta/y_1, \dots, y_n)$$

$$f(y|\mu) \propto e^{-\frac{1}{2\sigma^2}(y-\mu)^2} \tag{1}$$

Equation (1) is the kernel distribution for the normal density function.

Elicitation of the prior distribution

We have a prior distribution that is normal with mean m and variance, s^2 . Ignoring the part that doesn't involve μ , the shape of the prior density is given by

$$g(\mu) \propto e^{-\frac{1}{2s^2}(\mu-m)^2} \tag{2}$$

Posterior distribution for the Conjugate Prior

$$p(\mu|y) = g(\mu) \times f(y|\mu)$$

$$\propto e^{-\frac{1}{2}\left[\frac{(\mu-m)^2}{s^2} + \frac{(y-\mu)^2}{\sigma^2}\right]} \tag{3}$$

Putting the terms in exponent over the common denominator and expanding them out gives

$$\propto e^{-\frac{1}{2}\left[\frac{\sigma^2(\mu^2 - 2\mu m + m^2) + s^2(y^2 - 2y\mu + \mu^2)}{\sigma^2 s^2}\right]}$$

Combing the like terms

$$\propto e^{-\frac{1}{2}\left[\frac{(\sigma^2 + s^2)\mu^2 - 2(\sigma^2 m + s^2 y)\mu + m^2 \sigma^2 + y^2 s^2}{\sigma^2 s^2}\right]}$$

and factor out $(\sigma^2 + s^2)/\sigma^2 s^2$. Completing the square and absorbing the part that doesn't depend on μ into the proportionality constant, we have

$$\begin{aligned} &\propto e^{-\frac{1}{2\sigma^2s^2/(\sigma^2+s^2)}\left[\mu^2-2\frac{(\sigma^2m+s^2y)}{\sigma^2+s^2}\mu+\left(\frac{(\sigma^2m+s^2y)}{\sigma^2+s^2}\right)^2\right]} \\ &\propto e^{-\frac{1}{2\sigma^2s^2/(\sigma^2+s^2)}\left[\mu-\frac{(\sigma^2m+s^2y)}{\sigma^2+s^2}\right]^2} \end{aligned} \tag{4}$$

Thus, the posterior mean,

$$m' = \frac{(\sigma^2m + s^2y)}{\sigma^2 + s^2} \tag{5}$$

and

The posterior variance,

$$(s')^2 = \frac{(\sigma^2m + s^2y)}{\sigma^2 + s^2} \tag{6}$$

Hence, the prior has a *normal* (m, s²) and also, the posterior has a *normal* [m', (s')²] . This shows that the *normal* (m, s²) distribution is the conjugate family for the normal observation distribution with known variance.

Credible Interval

A credible interval is an interval that contains the probability of the parameter. It is an interval in the domain of a posterior probability distribution used for interval estimation.

$$= E(\mu|y) = \pm Z_{\frac{\alpha}{2}} SD(\mu|y) = \tag{7}$$

Where,

$E(\mu|y) ==$ *Posterior Mean*, $SD(\mu|y) ==$ *Posterior Standard Deviation*

and $Z_{\frac{\alpha}{2}} = z$ value

3. Data Analysis

Table 1 shows cross tabulation of sex and age. It is clearly seen that for each age group, women have more cases compare to men. The female average age was 53.10 while the male average age was 53.50. Also, the age group 52-61 years had the highest occurrence of hypertension.

Table 1: Data on the age and sex distribution for the hypertensive patients

Age Distribution	Sex		Total
	Female	Male	
2 - 11yr	5	5	10
12 - 21yr	4	0	4
22 - 31yr	24	18	42
32 - 41yr	134	29	163
42 - 51yr	227	72	299
52 - 61yr	252	90	342
62 - 71yr	151	86	237
72 - 81yr	52	10	62
82 - 91yr	12	2	14
92yr+	1	0	1
Total	862	312	1174

Determination of Distribution of the Data Using Normal Curve

From the figure 1 below, it is observed that the data is normally distributed. Therefore, normal distribution density function will be employed to obtain the data likelihood.




Figure 1: Normal Curve Depicting Data Distribution

Therefore,

$$\mu = 53.23 \text{ and } \sigma = 13.168$$

The mean and standard deviation of the data are 53.23 and 13.168 respectively.

Testing for the goodness of fit

H_0 = The distribution of the age follows normal distribution

H_1 = The distribution of the age does not follow normal distribution.

Table 2 shows that normal distribution was rank the best among other distribution that was in line with the data and was accepted at 98% and 99% Confidence Interval.

Table 2: Chi-square Easy-fit Output

Chi-Squared					
Deg. of freedom	10				
Statistic	19.312				
P-Value	0.03648				
Rank	1				
α	0.2	0.1	0.05	0.02	0.01
Critical Value	13.442	15.987	18.307	21.161	23.209
Reject?	Yes	Yes	Yes	No	No

Bayesian Analysis

A non-informative and informative prior will be used to ascertain the average age of hypertensive patients in the area under coverage.

Posterior Mean and Standard Deviation Using Uniform Prior

The Bayes Estimate (Posterior mean) of the age distribution of the patients is;

$$E(\mu|y) = \sum_{i=1}^{1174} \mu \times P(\mu|y) = 53.40156$$

And the Posterior Variance is;

$$V(\mu|y) = E(\mu^2|y) - [E(\mu|y)]^2$$

Where,
$$E(\mu^2|y) = \sum_{i=1}^{1174} \mu^2 \times P(\mu|y) = 2934.494$$

Therefore,

$$V(\mu|y) = 2934.494 - (53.40156)^2$$

$$V(\mu|y) = 82.76787$$

Also,

$$\text{Posterior Standard Deviation} = \sqrt{V(\mu|y)}$$

$$= \sqrt{82.76787}$$

$$\underline{\underline{= 9.097685}}$$

Posterior Mean and Standard Deviation Using Conjugate Prior

Using the conjugate prior of a Normal Distribution for a Normal Distribution likelihood with mean and variance parameters. It is given as:

$$f(y|\mu) = e^{-\frac{1}{2\sigma^2}(y-\mu)^2}$$

The Posterior Mean (Bayes Estimate) of θ is

$$E(\mu|y) = m' = \frac{(\sigma^2 m + s^2 y)}{\sigma^2 + s^2} = 53.7014$$

Where Posterior Variance is

$$V(\mu|y) = (s')^2 = \frac{(\sigma^2 m + s^2 y)}{\sigma^2 + s^2} = 10.8921$$

Therefore, the Posterior Standard Deviation is = 3.3003

The credible interval

$$= \text{Posterior mean of point estimate} \pm Z_{\alpha/2} \sqrt{(\text{Posterior Precision})}$$

$$= E(\mu|y) \pm Z_{\frac{\alpha}{2}} SD(\mu|y)$$

Where,

$$E(\mu|y) = 53.7014, \quad SD(\mu|y) = 3.3003 \quad \& \quad Z_{\frac{\alpha}{2}} = 1.96$$

Therefore,

The 95% credible interval for $(\mu|y)$ {the average age of hypertensive patients}: [47.23, 60.17]

4. Results and Summary

A total number of 1174 patients' were used for the analysis. The patients' ages ranged from 2 to 99 years which were grouped into 10 classes each of size 10. Most of the patients [342(29.1%)] fall within the age group 52 and 61 years while only 1 patient falls within the age group 92 and 99 years. The average age of the patients is $53\text{years} \pm 13.4568$. 862(73.4%) of the patients were female while 312(26.6%) of them were male. The female average age was 53.10 ± 13.203 while the male average age was 53.50 ± 13.301 . Chi-square and normal curve were used to show the age distribution of the respondents follow normal distribution. Using a positive uniform prior, the posterior follows a normal distribution. The Bayes estimate of the age distribution of hypertensive patients was 53.402 while its standard deviation was 9.098. Using a Conjugate prior, the likelihood and the posterior follow a normal distribution. The Bayes estimate of the age distribution of hypertensive patients was 53.7014 while its standard deviation was 3.3003. And the posterior confidence interval for average age of hypertensive patients is [47.23, 60.17].

5. Conclusion

This research outlines the basic goal of Bayesian inference, which is understood as a weighted average between knowledge about the parameter before data is observed and the information about the parameters (average age of hypertensive patients) contained in the observed. In all, the following findings were made: Having grouped data into 10 age groups, the frequency distribution of age groups of hypertensive patients in the area under study was obtained and the age group that had the highest occurrence is 52-61 years. Age distribution of hypertensive patients in the area under study follows a normal distribution. The average age of hypertensive patients in the area was obtained to be 53.23. For the classical approach, non-informative and informative prior give the same average age (53 years) but 13.168, 9.098 and 3.3003 standard deviations respectively. Credible Interval for posterior mean obtained was [47.23, 60.17]

References

- [1] Amoah AGB. Hypertension in Ghana: a cross-sectional community prevalence study in Greater Accra. *Ethn Dis.*, 13(2003):310-315.

- [2] Bartosh SM, Aronson AJ. Childhood hypertension: an update on etiology, diagnosis, and treatment. *Pediatr Clin North Am.*, 46(2)(1999): 235-252.
- [3] Bradshaw D, Groenewald P, Laubscher R, Nannan N, Nojilana B et al.; Initial burden of disease estimates for South Africa, 2000. *South African Medical Journal*, 93(9)(2003)
- [4] Delbecq A, Van de Ven A, Gustafson D. *Group techniques for program planning: A guide to nominal group and delphi process*. Glenview, IL: Scott, Foresman,1975.
- [5] European Society of Hypertension/European Society of Cardiology. Guidelines for the management of arterial hypertension. *Journal of Hypertension*, 21(2003):1011-1053.
- [6] Franklin SS, Gustin W, Wong ND, Larson MG, Weber MA, Kannel WB, et al. Hemodynamic patterns of age-related changes in blood pressure. The Framingham Heart Study. *Circulation*, 96(1997): 308-15.
- [7] Gelman, A., Carlin, J. B., Stern, H. S. and Rubin D. B.; *Test in Statistical Science: Bayesian Data Analysis*. Second Edition, (2004):
- [8] González-Velázquez F, Mendez GF. Cardiovascular risk stratification by means of the SCORE system in healthcare workers in Veracruz, Mexico. *Int J Cardiol*. 121(2006):81-83.
- [9] Hillege HL et al. Urinary albumin excretion predicts cardiovascular and noncardiovascular mortality in general population. *Circulation*, 106(2002):1777-1782.
- [10] JNC 6. National High Blood Pressure Education Program. The sixth report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Arch Intern Med.*, 157(1997): 2413-46.
- [11] Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: Analysis of worldwide data. *Lancet*, 365(2005):217-23.
- [12] L, Rodicio JL, eds. *Hypertension*. London, McGraw Hill International, 2001: 63-71.
- [13] Last JM, Abramson JH. *A dictionary of epidemiology*. 3rd ed. New York, NY: Oxford University Press, (1995).
- [14] Mancia G et al. Ambulatory blood pressure is superior to clinic blood pressure in predicting treatment induced regression of left ventricular hypertrophy. *Circulation*, 95(1997): 1464-1470.
- [15] Mayosi B, Fisher A, Lalloo U, Sitas F, Tollman S & Bradshaw D. Health in South Africa 4. The Burden of non-communicable diseases in South Africa. *The Lancet* 374(9693) (2009): 934- 947.
- [16] Messerli FH et al. Serum uric acid in essential hypertension: an indicator of renal vascular involvement. *Annals of Internal Medicine*, 93(1980): 817-821.
- [17] Minematsu K, Omae T.; Detection of damage to the brain. In: Zanchetti A, Hansson Mogensen CE.; Microalbuminuria and hypertension with focus on type 1 and 2 diabetes. *Journal of Internal Medicine*, 254(2003): 45-66.

- [18] Murray CJ, Lopez AD. Mortality by cause for eight regions of the world. Global Burden of Disease. *Lancet*. 349(1997):1269-1276.
- [19] National Heart, Lung, and Blood Institute. Report of the Second Task Force on Blood Pressure Control in Children. *Pediatrics*, 79(1)(1987): 1-25.
- [20] National High Blood Pressure Education Program. Report of the National High Blood Pickering T. Recommendations for the use of home (self) and ambulatory blood pressure monitoring. American Society of Hypertension Ad Hoc Panel. *American Journal of Hypertension*, 9(1996): 1-11.
- [21] Pressure Education Program Working Group on High Blood Pressure in Pregnancy. *Am J Obstet Gynecol*, 183(2000): S1-22.
- [22] Roccella E, Kaplan N. Interpretation and evaluation of clinical guidelines. In: Izzo JL Jr, Black HR (editors): *Hypertension Primer: The Essentials of High Blood Pressure: Basic Science, Population Science, and Clinical Management*. Philadelphia, PA: Lippincott Williams & Wilkins; 2003. pp. 126-7.
- [23] Ruilope LM et al. (2001). Renal function and intensive lowering of blood pressure in hypertensive participants of the Hypertension Optimal Treatment (HOT) study. *Journal of the American Society of Nephrology*, 12(2001): 218-225.
- [24] Sakuma M et al.; Reproducibility of home blood pressure measurements over a 1-year period. *American Journal of Hypertension*, 10(1997): 798-803.
- [25] Shaughnessy AF, Slawson DC, Bennett JH.; Becoming an information master: A guidebook to the medical information jungle. *J Fam Pract*, 39(1994): c489-99.
- [26] Sheps SG, Roccella EJ.; Reflections on the sixth report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Curr Hypertens Rep* 1(1999): 342-5.
- [27] Skaal L and Pengpid S.; Obesity and health problems among South African healthcare workers: do healthcare workers take care of themselves? *S Afr Fam Pract* . 53(6)(2011): 563-567.
- [28] Slawson DC, Shaughnessy AF.; Obtaining useful information from expert based sources. *BMJ*, 314(1997): 947-9.
- [29] Steyn NP, Bradshaw D, Norman R, et al. (2006). Dietary changes and the health transition in South Africa 2006: implications for health policy. Cape Town: *South African Medical Research Council*.
- [30] The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: The JNC 7 Report. *JAMA*, 289(2003): 2560-72.
- [31] U.S. Department of Health and Human Services, National Heart, Lung, and Blood Institute. 2001.

- [32] Vasani RS, Beiser A, Seshadri S, Larson MG, Kannel WB, D'Agostino RB, et al.; Residual lifetime risk for developing hypertension in middle-aged women and men: The Framingham Heart Study. *JAMA*, 287(2002):1003-10.
- [33] Vasani RS, Larson MG, Leip EP, Kannel WB, Levy D. Assessment of frequency of progression to hypertension in nonhypertensive participants in the Framingham Heart Study: A cohort study. *Lancet*, 358(2001):1682-6.
- [34] WHO (2011) *Global status report on non-communicable diseases*. Geneva: WHO
- [35] WHO (2011) Non Communicable Diseases country profile. Geneva: WHO.
- [36] WHO (2011) *Report on the Global Tobacco Epidemic: Warning about the dangers of tobacco*. Geneva.
<http://whqlibdoc.who.int/publications/2011/9789240687813> (accessed 14 May 2013)
- [37] WHO (2011) *Serum retinol concentrations for determining the prevalence of vitamin A deficiency in populations*. Vitamin and Mineral Nutrition Information System. Geneva: WHO.
<http://www.who.int/vmnis/indicators/retinol.pdf>
- [38] WHO (2013) Global Health Observatory Data Repository.
<http://apps.who.int/gho/data/node.main.A1079>
- [39] WHO African Region (2013). *Stakeholders' Dialogue to Address Risk Factors for Non-communicable Diseases in the WHO African Region*. Meeting held in Johannesburg, South Africa
- [40] Wolf-Maier K, Cooper RS, Banegas JR, Giampaoli S, Hense H, Joffres M, Katarinen M, Poulter N, Primatesta P, Rodriguez-Artalejo F, Stegmayr B, Thamm M, Tuomilehto J, Vanuzzo D, Vescio F.; Hypertension prevalence and blood pressure levels in 6 European countries, Canada, and the United States. *JAMA*. 289(2003): 2363-2369.
- [41] Whelton PK, He J, Appel LJ, Cutler JA, Havas S, Kotchen TA, et al.; Primary prevention of hypertension: Clinical and public health advisory from The National High Blood Pressure Education Program. *JAMA*; 288(2002):1882-8
- [42] William M. Bolstad, *Introduction to Bayesian Statistics*, John Wiley and Sons Inc. Hoboken, New Jersey, 2004 Edition.